

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

STIC-ILL

1545094

107/18

From: Baker, Maurie
Sent: Wednesday, July 17, 2002 6:30 PM
To: STIC-ILL
Subject: Please provide reference ASAP

404347

Please provide the following references ...ASAP
My mailbox is now located in *** 3B19 ***
Thank you!

TITLE: EFFECT OF NEW BIOLOGICALLY ACTIVE POLYPEPTIDES ON DIHEXADECYL PHOSPHATE
VESICLES.
AUTHOR(S): KRAUS J-L; MENASSA P
CORPORATE SOURCE: UNIV. D'AIX-MARSEILLE II, FAC. SCI. LUMINY CASE 901, LAB. CHEMIE
BIOMOLECULAIRE, 13288
MARSEILLE CEDEX 9, FR.
SOURCE: PHARMACOL RES COMMUN, (1987) 19 (7), 469-478.
CODEN: PLRCAT. ISSN: 0031-6989.
FILE SEGMENT: BA; OLD
LANGUAGE: English

09/758957: POLYPHARMACOPHORES

Maurie Garcia Baker, Ph.D.
Patent Examiner, Art Unit 1627
Crystal Mall 1, Rm. 3D11 (703) 308-0065

3671436

EFFECT OF NEW BIOLOGICALLY ACTIVE POLYPEPTIDES

ON DIHEXADECYL PHOSPHATE VESICLES ¹

Jean-Louis KRAUS and Pierre MENASSA ²

Université d'Aix-Marseille II, Faculté des Sciences de Luminy Case 901
Laboratoire de Chimie Biomoléculaire
13288 MARSEILLE CEDEX 9 - FRANCE

Received in final form 21 June 1987

SUMMARY . Effect of new biologically active polypeptides on hexadecyl phosphate vesicles. - The effect that monomeric and trimeric chemotactic peptides may have on the permeation of DHP vesicles was studied by means of electronic absorption spectroscopy. The results show that attaching covalently two or more of the same chemotactic peptides to a suitable carrier molecule produced changes in the permeation of DHP vesicles that deviate from simple additivity rule. Thus a parallelism between the biological potency of the peptides and DHP permeation exists.

Dihexadecylphosphate, vesicles, chemotactic peptides, permeation.

¹ Abbreviations : DHP = dihexadecyl phosphate, $\text{Ru}(\text{bpy})_3^{2+}$ = tris(2,2'-bipyridyl) Ruthenium (II) chloride hexahydrate, f-MLP = N-formyl L-methionyl-L-Leucyl-L-phenylalanine, tBoc = tertbutoxycarbonyl, DMF = N,N-dimethylformamide

² Present address : Concordia University , chemistry department
1455 De Maisonneuve. W. MONTREAL.Qué. CANADA H3G 1M8

In the past decade the functional properties of membrane assemblies were the object of investigation by scientists belonging to the physical, chemical or biological community. These researches were triggered by the important role that plasma membranes played as a molecular barrier in biological cells.

Single compartment vesicles can be considered as a model for the study of membrane permeability. Indeed, like real membranes they are made from bilayers. In this work, single compartment vesicles made from DHP were used as a membrane model to ascertain a new concept in pharmacology. This concept based on the finding that attaching covalently two or more of the same pharmacophores (biologically active molecules) to a suitable carrier molecule may produce synergetic biological effects (KRAUS. 1984 and papers mentioned therein). Indeed it has been pointed out that polypharmacophores may interact with the receptors located on the cell surface causing them to cluster (STAHL. 1978 ; KAWASAKI. 1978 ; KORNFELD. 1976 ; LEE. 1978 ; DINTZIS. 1982 ; TOKES. 1982 ; POMPIPOM. 1981). This clustering effect of membrane receptors may induce changes in the chemico-physical properties of the membrane, among them permeability. The marker $\text{Ru}(\text{bpy})_3^{2+}$ was encapsulated in DHP vesicles. The permeation of DHP vesicles by various monomeric and polymeric peptides (scheme I) derived from the well-known chemotactic peptide f-MLP (SHOWELL. 1976) was studied by monitoring $\text{Ru}(\text{bpy})_3^{2+}$ leakage across the bilayer using visible absorption spectroscopy.

MATERIALS AND METHODS

The unitary monomeric peptide analogs f-MetLeuPheOCH₃ 6 and BOC-PheLeuPheOCH₃ 7 were obtained from Armand Frappier Corporation (Mon-

of membrane assemblies belonging to the physical, ves were triggered by the as a molecular barrier in d as a model for the study ranes they are made from s made from DHP were used in pharmacology. This con- ly two or more of the same a suitable carrier mole- (KRAUS. 1984 and papers at that polypharmacophores ll surface causing them to ELD. 1976 ; LEE. 1978 ; his clustering effect of ico-physical properties of r $\text{Ru}(\text{bpy})_3^{2+}$ was encapsu- cles by various monomeric he well-known chemotactic monitoring $\text{Ru}(\text{bpy})_3^{2+}$ lea- spectroscopy.

f-MetLeuPheOCH₃ 6 and appier Corporation (Mon-

treal) Canada. L-Lys-L-Lys, 2 HCl 5 was purchased from Sigma. N,N'-bis (2-hydroxyethyl)-4,13 diaza-18 crown-6 3 been prepared according to Gatto and Gokel procedure (1984) . The coupling with N-tBOC MetLeuPheOH was accom- plished using dicyclohexylcarbodiimide in DMF in the presence of 1-hydroxy- benzotriazole as catalyst (KONIG. 1970) . Direct deblocking and formyla- tion of the terminal methionine residue was carried out as recently des- cribed (LAJOIE. 1984) leading to compound 1. Compound 2 was obtained by coupling 3 with N-tBOCPheLeuPheOH . Condensation of the p-nitrophenylester of N-tBOCMetLeuPheOH with the L-Lysyl-L-Lysyl trifluoroacetate salt in DMF gave a high yield of the corresponding N-tBOC protected intermediate. The latter was deblocked and formylated as described above, leading to compound 4.

All products thus obtained were chromatographically pure as judged by thin layer chromatography and their structures were established by IR, NMR (200 MHz) and mass spectrometry. Satisfactory elemental analyses were obtained in all cases.

Biological activity of these peptides analogs was evaluated using the release of lysozyme from human neutrophils as the assay method (SHOWELL. 1976) . Reproducible results were readily obtained using cells from heal- thy blood donors. The lysosomal enzyme inducing activity of each peptide derivative was computed from dose-response curves to give the ED50 values (the concentration of compound causing 50% of the maximal release of lyso- zyme).

DHP vesicles were prepared by sonication with an ultrasonics cell Disruptor filled with a microtip set at 35 W of output power. Typically, 27 mg of DHP was first heated to 80°, then 33 ml of distilled water pre-heated

at 80°C were added, giving a DHP concentration $1.5 \cdot 10^{-3} \text{ mol.l}^{-1}$. After initial sonication 0.1 mol.l^{-1} NaOH was injected to give an NaOH concentration of $7.5 \cdot 10^{-4} \text{ mol.l}^{-1}$. For entrapment purposes appropriate volume of Ru(bpy)_3^{2+} were injected after a few minutes of initial sonication (TRICOT, 1983). Sonication was then continued until a constant turbidity was reached. Traces of titanium released by the microtip during sonication were removed by centrifugation. Externally adsorbed cations, those entrapped during sonication but able to readily permeate the vesicle bilayer, and those in the outer bulk solution, could be removed by passing the vesicle dispersion through a column of Bio-Rad AG 50W-X2 cation exchange resin (100-200 mesh, hydrogen form).

Levels of entrapment were determined at room temperature from the residual concentration of Ru(bpy)_3^{2+} measured after treatment with the cation exchange resin. Concentrations of Ru(bpy)_3^{2+} were determined spectrophotometrically at 454 nm (TRICOT et al. 1984). The absorbance/scattering spectrum of the vesicle dispersion in the absence of Ru(bpy)_3^{2+} was determined for each sample and corrections made when determining concentrations of Ru(bpy)_3^{2+} from spectrometric measurements. At $1.5 \cdot 10^{-3} \text{ mol.l}^{-1}$ DHP, as used in entrapment experiments, no destabilization of vesicles due to pH changes during resin treatment was observed. (TRICOT, FURLONG, MAU and SADE, 1985).

Before further use of the vesicles with entrapped ions, HCl resulting from the cation exchange resin was removed by ultrafiltration.

Permeability values are reported as PD50's values. PD50 is the required amount of pharmacophores needed to induce at 50% the permeability of DHP vesicles to Ru(bpy)_3^{2+} .

RESULTS AND DISCUSSION

The dimeric and trimeric peptides analogs in Scheme I were synthesized with a view to testing a possible relation between the potency of these compounds to induce lysosomal enzyme release and their effect on the permeation of DHP vesicles. The effect that monomeric analogs 6, 7 and the backbone 3, 5 may have on the permeation of DHP vesicles was also assessed.

The calculated PD50 of FMLP was found to be 4.5×10^{-3} M.L⁻¹. The PD50's of the dimeric 1 peptide and that of the trimeric peptide 4 were found to be 6.3×10^{-5} M.L⁻¹ and 6.0×10^{-5} M.L⁻¹ respectively. This clearly shows that the trimeric and the dimeric species permeate DHP vesicles by orders to magnitude more than the monomeric species. The PD50's of the carrier molecules were also calculated. The PD50 of Lys-Lys, 2 HCl 5 was found to be 4.15×10^{-4} M.L⁻¹, and 8.10×10^{-4} M.L⁻¹ for compound 3. Thus the carrier molecule by itself has a noticeable effect. This can be understood at least for 3, which can be considered as an ionophore. However this effect alone cannot explain the large difference between on the one hand the PD50 of the monomeric species and on the other hand those of the dimeric and trimeric species. Therefore the observed synergetic induced permeation is in part due to the carrier molecules and in part due to the covalent attachment of two or three FMLP entities to those carrier molecules. The results obtained are in line with the observed biological effects (induction of lysosomal enzyme release from human neutrophils). Table I regroups the PD50's of the different studied species as well as their ED50's. Two more compounds known as lysosomes release antagonists (DAY, 1980) were also studied BOC-Phe-Leu-Phe-OCH₃ 7 and its dimer analog 2.

SCHEME 1







TABLE I

Comparison between the PD50's of the different species studied.
The ED50's of some chemotactic peptides are also included.

COMPOUNDS	Biological Results		Permeation Results	
	ED50 (Mol.L ⁻¹) Lysosyme release	Relative Potency %	PD50 (Mol.L ⁻¹)	Relative Potency %
$\underline{6}$	100×10^{-9}	100	4.5×10^{-3}	100
$\underline{1}$	2.0×10^{-9}	5000	6.3×10^{-5}	7000
$\underline{4}$	3.0×10^{-10}	33000	6.0×10^{-5}	7500
$\underline{3}$	no activity	-	8.1×10^{-4}	550
$\underline{5}$	no activity	-	4.15×10^{-4}	1100
$\underline{7}$	antagonist	-	2×10^{-3}	100
$\underline{2}^*$	antagonist	-	6.5×10^{-5}	3100

* The relative potency of $\underline{2}$ was calculated with respect to that of $\underline{7}$



cies studied.
ncluded.

Permeation Results

PD50 (Mol.L ⁻¹)	Relative Potency %
4.5 x 10 ⁻³	100
6.3 x 10 ⁻⁵	7000
6.0 x 10 ⁻⁵	7500
8.1 x 10 ⁻⁴	550
4.15 x 10 ⁻⁴	1100
2 x 10 ⁻³	100
6.5 x 10 ⁻⁵	3100

spect to that of 7

Their PD50's were found to be 2.0×10^{-3} M.L⁻¹ and 6.5×10^{-5} M.L⁻¹ respectively. This is in line with what one would have expected from the previous discussion.

The use of synthetic phospholipid such as DHP in the preparation of bilayered model membrane in place of natural phospholipids is probably not the ideal model of membrane, since the normal phospholipid components of the cell surface present on their outer layer all the possible chemical groups required to establish the interaction with the exogenous peptides but the permeation results show discrepancies which are in line with the lysosomal enzyme activity. Of course the intensity of the biological responses ED50 and PD50 cannot be directly and quantitatively correlated since polypharmacophoric compounds 4 (ED50 33000 at 3.10^{-10} M) and compound 1 (ED 50 5000 at 2.10^{-9} M) have practically the same permeation activity but in comparison with the corresponding monomeric compound 6 the PD 50 values are significantly different almost two order magnitude.

ACKNOWLEDGEMENTS

Gratitude is expressed to Dr. B. Belleau from Mc Gill University and to professor C. Sandorfy from the Université de Montréal for providing facilities, and for stimulating discussions. We thank Miss Antonia di Paola for biological testing. We (JLK) gratefully acknowledge the France-Québec exchange program for work at McGill University.

REFERENCES

- 1 - DAY, A.R., D. PINON, N. MUTHUKUMARASWAMY and R.J. FREER. Peptides, 4 : 289 - 291, 1980.
- 2 - DINTZIS, R.Z., B. VOGELSTEIN, and H.M. DINTZIS. Proc. Natl. Acad. Sci. USA. 79 : 884 - 888, 1982.

- 3 - GATTO, V.J. and G.W. GOKEL. J. Amer. Chem. Soc., 106 : 8240 - 8244, 1984.
- 4 - KAWASAKI, T., R. ETOH and I. YAMASHINO. Biochem. Biophys. Res. Commun. 81 : 1018 - 1024 , 1978.
- 5 - KRAUS, J.L., A. DI PAOLA, and B. BELLEAU. Biochem. Biophys. Res. Commun. 124 : 939 - 944, 1984.
- 6 - KONIG, W. and R. GEIGER. Chem. Ber. 103 : 788 - 798, 1970.
- 7 - KORNFELD, R., and S. KORNFELD. Annu. Rev. Biochem. 45 : 217 - 237, 1976.
- 8 - LAJOIE, G. and J.L. KRAUS. Peptides. 5 : 653 - 654, 1984.
- 9 - LEE, Y.C. Carbohydr. Res. 67 : 509 - 514, 1978.
- 10 - POMPIPOM, M.M., R.L. BUGNASI, J.C. ROBBINS, T.W. DOEBBER and T.Y. SHEN. J. Med. Chem. 24 : 1388 - 1395, 1981.
- 11 - SHOWELL, H.J., R.J. FREER, S.H. ZIGMOND, E. SCHIFFMANN, S. ASWANI-KUMAR, B. CORCORAN, and E.L. BECKER. J. Exp. Med. 143 : 1154 - 1169, 1976.
- 12 - STAHL, P.D., J.S. RODMAN, J.S. MILLER, and P.H. SCHLESINGER. Proc. Natl. Acad. Sci. USA. 75 : 1399 - 1403, 1978.
- 13 - TOKES, Z.A., K.E. ROGERS and A. REMBAUM. Proc. Natl. Acad. Sci. USA. 79 : 2026 - 2030, 1982.
- 14 - TRICOT, Y.M. , FURLONG, D.N., SASSE, W.H.F., DAIVIS, P., and SNOOK, I. Aust. J. Chem. 36 : 609 -612, 1983.
- 15 - TRICOT, Y.M., FURLONG, D.N., SASSE, W.D.F., DAIVIS, P., SNOOK, I and VAN HEGEN, W. J. Colloid Interface Sci., 97, 380-391, 1984.
- 16 - TRICOT, Y.M., FURLONG, D.N., and SASSE W.H.F. Aust. J. Chem., 37 , 1147-56 , 1984.

oc., 106 : 8240 - 8244,

m. Biophys. Res. Commun.

Biochem. Biophys. Res.

798, 1970.

iochem. 45 : 217 - 237,

654, 1984.

3.

4. DOEBBER and T.Y. SHEN.

SCHIFFMANN, S. ASWANI-

Med. 143 : 1154 - 1169,

P.H. SCHLESINGER. Proc.

c. Natl. Acad. Sci. USA.

AIVIS, P., and SNOOK, I.

AIVIS, P., SNOOK, I and

180-391, 1984.

F. Aust. J. Chem., 37 ,

17 - TRICOT, Y.M., FURLONG, D.N., MAU, A.W. and SASSE, W.H.S. Aust. J.
Chem., 38, 527-535, 1985.

=> d his

(FILE 'HOME' ENTERED AT 16:37:32 ON 17 JUL 2002)

FILE 'REGISTRY' ENTERED AT 16:37:51 ON 17 JUL 2002

L1 STRUCTURE UPLOADED
L2 0 S L1
L3 STRUCTURE UPLOADED
L4 0 S L3
L5 7 S L3 FULL

FILE 'CAOLD' ENTERED AT 16:41:47 ON 17 JUL 2002

L6 3 S L5

FILE 'STNGUIDE' ENTERED AT 16:43:38 ON 17 JUL 2002

FILE 'CAPLUS, MEDLINE, BIOSIS, SCISEARCH, EMBASE, USPATFULL' ENTERED AT
16:47:33 ON 17 JUL 2002

L7 3470 S POLYPHARMA?
L8 2 S POLYPHARMACOPHORE
L9 4 S POLYPHARMACOPHO?
L10 2 S L9 NOT L8

FILE 'STNGUIDE' ENTERED AT 16:51:49 ON 17 JUL 2002

FILE 'CAPLUS, MEDLINE, BIOSIS, SCISEARCH, EMBASE, USPATFULL' ENTERED AT
16:57:46 ON 17 JUL 2002

L11 6938 S DOPAMINE (P) TRANSPOR? (P) INHIBI?
L12 880 S L11 AND (POLY? OR DIME? OR TRIM?)
L13 537 S L11 AND (DIME? OR TRIM?)
L14 138 S L11 AND (TRIM?)
L15 37 S L11 AND (TRIMER?)
L16 1258 S DOPAMINE (3A) TRANSPOR? (3A) INHIBI?
L17 1 S L16 AND (TRIMER?)
L18 112 S L16 AND (MULTI?)
L19 66 DUP REM L18 (46 DUPLICATES REMOVED)
L20 14 S L16 AND (DIMER OR DIMERIC OR BIVALENT OR DIVALENT OR LINKER)
L21 14 DUP REM L20 (0 DUPLICATES REMOVED)
L22 45 S L16 AND (LINKED)
L23 31 DUP REM L22 (14 DUPLICATES REMOVED)

=> log y